

# ON THE NUMBER OF MAXIMAL INDEPENDENT SETS IN A GRAPH

DAVID R. WOOD

ABSTRACT. Miller and Muller (1960) and independently Moon and Moser (1965) determined the maximum number of maximal independent sets in an  $n$ -vertex graph. We give a new and simple proof of this result.

Let  $G$  be a (simple, undirected, finite) graph. A set  $S \subseteq V(G)$  is *independent* if no edge of  $G$  has both its endpoints in  $S$ . An independent set  $S$  is *maximal* if no independent set of  $G$  properly contains  $S$ . Let  $\text{MIS}(G)$  be the set of all maximal independent sets in  $G$ . Miller and Muller [7] and Moon and Moser [8] independently proved that the maximum, taken over all  $n$ -vertex graphs  $G$ , of  $|\text{MIS}(G)|$  equals

$$g(n) := \begin{cases} 3^{n/3} & \text{if } n \equiv 0 \pmod{3} \\ 4 \cdot 3^{(n-4)/3} & \text{if } n \equiv 1 \pmod{3} \\ 2 \cdot 3^{(n-2)/3} & \text{if } n \equiv 2 \pmod{3} \end{cases}.$$

This result is important because  $g(n)$  bounds the time complexity of various algorithms that output all maximal independent sets [1–3, 5, 6, 9, 10]. Here we give a new and simple proof of this upper bound on  $|\text{MIS}(G)|$ .

**Theorem 1** ([7, 8]). *For every  $n$ -vertex graph  $G$ ,*

$$|\text{MIS}(G)| \leq g(n) \text{ .}$$

*Proof.* We proceed by induction on  $n$ . The base case with  $n \leq 2$  is easily verified. Now assume that  $n \geq 3$ . Let  $G$  be a graph with  $n$  vertices. Let  $d$  be the minimum degree of  $G$ . Let  $v$  be a vertex of degree  $d$  in  $G$ . Let  $N[v]$  be the closed neighbourhood of  $v$ . If  $I \in \text{MIS}(G)$  then  $I \cap N[v] \neq \emptyset$ , otherwise  $I \cup \{v\}$  would

---

*Date:* January 19, 2013.

Department of Mathematics and Statistics, The University of Melbourne, Melbourne, Australia (woodd@unimelb.edu.au). Supported by a QEII Fellowship from the Australian Research Council.

be an independent set. Moreover, if  $w \in I \cap N[v]$  then  $I \setminus \{w\} \in \text{MIS}(G - N[w])$ . Thus

$$|\text{MIS}(G)| \leq \sum_{w \in N_G[v]} |\text{MIS}(G - N_G[w])| ,$$

Since  $\deg(w) \geq d$  and  $g$  is non-decreasing, by induction,

$$|\text{MIS}(G)| \leq (d+1) \cdot g(n-d-1) .$$

Note that

$$4 \cdot 3^{(n-4)/3} \leq g(n) \leq 3^{n/3} .$$

If  $d \geq 3$  then

$$|\text{MIS}(G)| \leq (d+1) \cdot 3^{(n-d-1)/3} \leq 4 \cdot 3^{(n-4)/3} \leq g(n) .$$

If  $d = 2$  then

$$|\text{MIS}(G)| \leq 3 \cdot g(n-3) = g(n) .$$

If  $d = 1$  and  $n \equiv 1 \pmod{3}$  then since  $n-2 \equiv 2 \pmod{3}$ ,

$$|\text{MIS}(G)| \leq 2 \cdot g(n-2) \leq 2 \cdot 2 \cdot 3^{(n-2-2)/3} = 4 \cdot 3^{(n-4)/3} = g(n) .$$

If  $d = 1$  and  $n \equiv 0 \pmod{3}$  then since  $n-2 \equiv 1 \pmod{3}$ ,

$$|\text{MIS}(G)| \leq 2 \cdot g(n-2) \leq 2 \cdot 4 \cdot 3^{(n-2-4)/3} < 3^{n/3} = g(n) .$$

If  $d = 1$  and  $n \equiv 2 \pmod{3}$  then since  $n-2 \equiv 0 \pmod{3}$ ,

$$|\text{MIS}(G)| \leq 2 \cdot g(n-2) \leq 2 \cdot 3^{(n-2)/3} = g(n) .$$

This proves that  $|\text{MIS}(G)| \leq g(n)$ , as desired.  $\square$

For completeness we describe the example by Miller and Muller [7] and Moon and Moser [8] that proves that Theorem 1 is best possible. If  $n \equiv 0 \pmod{3}$  then let  $M_n$  be the disjoint union of  $\frac{n}{3}$  copies of  $K_3$ . If  $n \equiv 1 \pmod{3}$  then let  $M_n$  be the disjoint union of  $K_4$  and  $\frac{n-4}{3}$  copies of  $K_3$ . If  $n \equiv 2 \pmod{3}$  then let  $M_n$  be the disjoint union of  $K_2$  and  $\frac{n-2}{3}$  copies of  $K_3$ . Observe that  $|\text{MIS}(M_n)| = g(n)$ .

Note that Vatter [11] independently proved Theorem 1, and gave a connection between this result and the question, “What is the largest integer that is the product of positive integers with sum  $n$ ?” Also note that Dieter Kratsch proved that  $|\text{MIS}(G)| \leq 3^{n/3}$  using a similar proof to that presented here; see Gaspers [4, page 177]. Thanks to the authors of [4, 11] for pointing out these references.

## REFERENCES

- [1] COEN BRON AND JOEP KERBOSCH. Finding all cliques of an undirected graph. *Comm. ACM*, 16(9):575–577, 1973. doi:10.1145/362342.362367.
- [2] DAVID EPPSTEIN. Small maximal independent sets and faster exact graph coloring. *J. Graph Algorithms Appl.*, 7(2):131–140, 2003. <http://jgaa.info/accepted/2003/Eppstein2003.7.2.pdf>.
- [3] DAVID EPPSTEIN, MAARTEN LÖFFLER, AND DARREN STRASH. Listing all maximal cliques in sparse graphs in near-optimal time. 2010. <http://arxiv.org/abs/1006.5440>.
- [4] SERGE GASPERS. *Exponential Time Algorithms: Structures, Measures, and Bounds*. VDM Verlag Dr. Müller, 2010. [http://www.kr.tuwien.ac.at/drm/gaspers/SergeBookETA2010\\_screen.pdf](http://www.kr.tuwien.ac.at/drm/gaspers/SergeBookETA2010_screen.pdf).
- [5] DAVID S. JOHNSON, MIHALIS YANNAKAKIS, AND CHRISTOS H. PAPADIMITRIOU. On generating all maximal independent sets. *Inform. Process. Lett.*, 27(3):119–123, 1988. doi:10.1016/0020-0190(88)90065-8.
- [6] EUGENE L. LAWLER, JAN KAREL LENSTRA, AND ALEXANDER H. G. RINNOOY KAN. Generating all maximal independent sets: NP-hardness and polynomial-time algorithms. *SIAM J. Comput.*, 9(3):558–565, 1980. doi:10.1137/0209042.
- [7] R. E. MILLER AND D. E. MULLER. A problem of maximum consistent subsets, 1960. IBM Research Report RC-240, J. T. Watson Research Center, New York, USA.
- [8] JOHN W. MOON AND LEO MOSER. On cliques in graphs. *Israel J. Math.*, 3:23–28, 1965. doi:10.1007/BF02760024.
- [9] ETSUJI TOMITA, AKIRA TANAKA, AND HARUHISA TAKAHASHI. The worst-case time complexity for generating all maximal cliques and computational experiments. *Theoret. Comput. Sci.*, 363(1):28–42, 2006. doi:10.1016/j.tcs.2006.06.015.
- [10] SHUJI TSUKIYAMA, MIKIO IDE, HIROMU ARIYOSHI, AND ISAO SHIRAKAWA. A new algorithm for generating all the maximal independent sets. *SIAM J. Comput.*, 6(3):505–517, 1977. doi:10.1137/0206036.
- [11] VINCENT VATTER. Maximal independent sets and separating covers. *American Mathematical Monthly*, 118:418–423, 2011. <http://www.math.ufl.edu/~vatter/publications/sss-mis/>.